MEMORANDUM

To: USACE Colonel James L. Booth, LTC Todd F. Polk, Richard McMillen, SFWMD Governing Board, Executive Director Drew Bartlett, Jennifer Reynolds, Lawrence Glenn, DEP Secretary Shawn Hamilton

From: Periodic Scientists Conference Call Participants

Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex

Holly Milbrandt & Dana Dettmar - City of Sanibel

Lesli Haynes & Lisa Kreiger - Lee County

Harry Phillips & Maya Robert - City of Cape Coral

Leah Reidenbach, Rick Bartleson PhD, & Matt Depaolis - SCCF (Sanibel-Captiva Conservation Foundation)

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: January 23 - 29, 2024

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

Caloosahatchee Conditions Summary: Flow to the Caloosahatchee Estuary had a 7-day average of 2,019 cfs at S-79 with a 7-day average of 917 cfs (45%) coming from the lake at S-77. The 14-day moving average flow at S-79 is 2,542 cfs and has been in the stress flow envelope (2,100 – 2,600 cfs; RECOVER 2020) for 15 days.

Recommendation: The prolonged high lake stage is having long-term negative impacts on the health of Lake Okeechobee. With El Niño conditions currently bringing above average rainfall this dry season, lowering the lake prior to the 2024 rainy season will prove to be challenging. Time is of the essence; we encourage the Corps to start managing Lake Okeechobee to reduce lake levels while maintaining optimum flows and use all available outlets to prevent damaging discharges to the estuaries.

USACE Action: With Lake Okeechobee stage in the Intermediate Sub-band, the Tributary Hydrologic conditions in the Wet category, Part D of the 2008 LORS suggests up to 4,000 cfs at S-77 and 1,800 cfs at S-80. On 6/10/23 the USACE increased releases from Lake Okeechobee to the Caloosahatchee Estuary from the W.P. Franklin Lock and Dam (S-79) to 2,000 cfs. Releases to the St. Lucie Estuary (S-80) remain at 0 cfs.

Lake Flows: In the past 7 days the total outflow from Lake Okeechobee was 19,178 AF with 12,733 AF to the Caloosahatchee through S-77, 59 AF to the St. Lucie canal though S-308, 40 AF through S-310 in Clewiston, 1,359 AF though the L8 canal, and 4,987 AF to the EAA through S-351, S-352, and S-354. The total net inflow to the Lake was 52,078 AF (52074 AF from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1) and a total backflow of 4 AF from the S-310. Water conservation areas received flows of 6,553 AF, 10,000 AF, and 11,304 AF at WCA1, WCA2, and WCA3, respectively. Everglades National Park received 11,390 AF.


Lake Level: 16.33 ft (Intermediate Sub-Band) Last Week: 16.17 ft Last Year: 16.04 ft

7-Day Lake Recession Rate: +0.16 ft/week


Lake Okeechobee Inflow: 3,978 cfs

Lake Okeechobee Outflow: 1,373 cfs

Weekly Rainfall Total: WP Franklin: 0.00" Ortona: 0.02" Moore Haven: 0.15"



ACOE Daily Reports					
Date	S79 Flow (cfs)				
1/23/24	2399	662	731		
1/24/24	2111	751	922		
1/25/24	1704	401	679		
1/26/24	1445	395	485		
1/27/24	1651	588	704		
1/28/24	2239	981	997		
1/29/24	2585	1628	1904		
7-day avg	2019	772	917		



Lio	ht	Pen	etr	ation	
	,			uuvii	

Site	25% lz	Target Values	Turbidity	Target Values	
	meters		NTU		
Fort Myers	0.6	> 1	2.5	< 18	
Shell Point	ND	>2.2	ND	< 18	
Causeway	2.7	> 2.2	2.3	< 5	

25% Iz is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

Cyanobacteria Status: On 1/30/24 sampling for cyanobacteria by the Lee County Environmental Lab reported the **presence** of *Microcystis* and *Dolichospermum* at the **Davis Boat Ramp** as some streaks. No potentially toxic species were found in sub surface samples at Moore Haven or Alva on 1/25/24.

Upper Estuary Conditions: The 30-day average surface salinity at the Fort Myers Yacht Basin was 1.4 psu, within the suitable range for tape grass.

Lower Estuary Conditions: The average salinity at Shell Point RECON was 20 psu, in the optimal range for oysters but below optimal for seagrass.

Water Quality Conditions:

Monitor Site	Salinity (psu) ^a [previous week]	Diss O ₂ (mg/L) ^b	FDOM (qsde) ^c	Chlorophyll (µg/L) ^d	Temperature (°F)
Beautiful Island	0.2 - 0.2 [0.2 - 0.2]	6.0 - 7.6	164 – 225	6.0	65.1 – 78.1
Fort Myers Yacht Basin	0.2 - 1.6 [0.2 - 0.5]	7.3 – 9.0	175 – 235	6.6	61.9 – 74.4
Shell Point	5.6 – 31 [3.3 – 29]	6.5 - 8.5			60.8 – 73.0
McIntyre Creek	20.6 - 28.6 [19.7 - 28.4]	2.9 - 9.2			61.4 – 74.2
Tarpon Bay	21.7 - 31.4 [16.8 - 28.6]	6.2 - 8.5	30.5 - 85.3	1.3 – 2.5	61.3 – 73.3
Wulfert Flats	21.5 – 31.0 [22.1 – 28.6]	6.3 - 9.7		4.9 – 16.2	61.3 – 73.4

Red values are outside of the preferred range.

Wildlife Impacts: In the past week, the CROW wildlife hospital on Sanibel admitted 9 patients with suspected red tide/toxicosis: 1 adult herring gull (still at CROW), 1 adult double-crested cormorant (deceased), 1 adult osprey (deceased), 1 adult white pelican (deceased) and 5 adult royal terns (1 still at CROW, 4 deceased).

Red Tide: On 1/26/24, the FWC reported that the red tide organism *Karenia brevis* was observed at background concentrations in **one sample** from **Pinellas County**.

Shellfish Advisory: Shellfish harvest area #6222/6232 Matlacha Pass Shellfish Harvest Area is OPEN by the Florida Department of Agriculture and Consumer Services (FDACS) as of 1/27/2024 because fecal coliform results indicate that water quality meets NSSP standards as defined in Chapter 5L-1.003, Florida Administrative Code

^a Salinity target values: BI < 5, FM < 10, SP = 10 - 30


b Dissolved O2 target values: all sites > 4

^c FDOM target values: BI < 70, FM < 70, SP < 11

^d Chlorophyll target values: BI < 11, FM < 11, SP < 11

s Single sonde lower and surface layer or surface grab lab measurement

⁻⁻⁻⁻⁻ no data



Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (Vallisneria americana) health (Ft. Myers only) conditions.

*Ft. Myers sensor is in the lower strata



Water clarity at Lighthouse Beach Park on 1/30/24 at 12:55 PM on a rising tide (1.2 ft). <u>Lighthouse Beach Park Virtual Tour.</u>