

MEMORANDUM

To: USACE Colonel Brandon L. Bowman, Major Cory Bell, Richard McMillen, SFWMD Governing Board,
Executive Director Drew Bartlett, Jennifer Reynolds, DEP Secretary Alexis Lambert

From: Periodic Scientists Conference Call Participants

Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex

Holly Milbrandt & Dana Dettmar - City of Sanibel

Allie Pecenka, Rick Bartleson PhD & Matt Depaolis- Sanibel-Captiva Conservation Foundation

With contributions from Harry Phillips & Maya Robert PhD- City of Cape Coral

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: **June 17- 23, 2025**

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

Caloosahatchee Conditions Summary: Flow to the Caloosahatchee Estuary had a 7-day average of **297 cfs** at **S-79** with a 7-day average of **85 cfs** (29%) coming from the lake at **S-77**. **The 14-day moving average flow at S-79 was 566 cfs** and has been **below the optimum flow envelope** (<750 cfs) for **4 days** after **15 days in the optimum flow envelope** (750- 2,100 cfs; RECOVER 2020). **The 14-day moving average flow at S-77 was 42 cfs.**

Recommendation: We request that flows to the Caloosahatchee at S-79 be within the optimum flow envelope to best support estuarine health during the onset of the wet season. Given the reduced flow schedule with a lake release target of 250 cfs and additional flows expected to be supplemented by basin runoff, we request that modeling continue to account for realized vs projected precipitation as it impacts S-79 flows. In the event basin runoff does not supplement S-79 flows to reach the optimum flow envelope, we request that additional releases from S-77 be made to achieve optimum flows of 750- 2,100 cfs at S-79.

USACE Action: Lake Okeechobee stage is in the lower portion of Zone D (Zone D3 of the PA25 simulation) of the LOSOM regulation schedule. ENSO-neutral conditions are present and are likely during the summer. The District will continue to monitor conditions in the estuaries, as well as the systemwide conditions, as the wet season progresses. With the initiation of the wet season, and the termination of Recovery Operations, Normal Lake Operations will resume pursuant to the considerations in LOSOM as informed by PA25. To maintain favorable salinity levels in the estuaries and begin to conserve water, it is recommended that flow targets for the Caloosahatchee Estuary should rely on rainy season basin flows to ensure the delivery of the Minimum Flow and Level, but use Lake Okeechobee flows from S-77 to ensure S-79 flows remain above a targeted steady release of 250 cfs; flow targets for the St. Lucie Estuary and Lake Worth Lagoon should remain at 0 cfs consistent with Normal Operations within Zone D. The District will continue to monitor salinity conditions in the estuaries and water supply conditions throughout the system as the wet season progresses to assess future operational recommendations.

Lake Flows: In the past 7 days the total outflow from Lake Okeechobee was **11,485 AF** with **1,191 AF** to the Caloosahatchee through **S-77**, **-1,783 AF** to the St. Lucie canal through **S-308** and **12,077 AF** to the EAA through **S-351**, **S-352**, and **S-354**. The total net inflow to the Lake was **3,709 AF** from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1. Water conservation areas received flows of **299 AF**, **2,493 AF**, and **9,094 AF** at **WCA1**, **WCA2**, and **WCA3**, respectively. Everglades National Park received **567 AF**.

*Data missing from S-310 and L-8 from 6/17- 6/23, from S-78 on 6/18 and 6/21 and from S-80 on 6/18.

Lake Level: 10.99 ft (Zone D3)

Last Week: 11.18 ft

Last Year: 13.39 ft

7-Day Lake Recession Rate: -0.19 ft/week

Lake Okeechobee Inflow: 318 cfs

Lake Okeechobee Outflow: 693 cfs


Weekly Rainfall Total: WP Franklin: 3.83"

Ortona: 2.65"

Julian Keen Jr.: 0.30"

Cyanobacteria Status: On 6/23/25, sampling for cyanobacteria by the Lee County Environmental Lab reported *Dolichospermum* and *Microcystis* as **present upstream of the Franklin Locks**, appearing as sparse streaks with slight accumulation and at the **Davis Boat Ramp**, appearing as sparse streaks.

Red Tide: On 6/20/25, the FWC reported that the red tide organism, *Karenia brevis*, was detected at **background concentrations** in one sample from **Southwest Florida**; from Lee County.



Light Penetration

Site	25% Iz		Turbidity	
	meters		NTU	
Beautiful Is	0.8	> 1	3.0	< 18
Shell Point	2.0	>2.2	1.8	< 18
Causeway	4.0	> 2.2	3.5	< 5

25% Iz is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

Upper Estuary Conditions: The 30-day moving average surface salinity at the Fort Myers Yacht Basin was **10 psu**, in the stress range for tape grass (10- 15 psu; RECOVER 2020) and has been **over 10.0 psu for 36 days**. The weekly average was 9.2 psu.

Lower Estuary Conditions: The weekly average salinity at the Shell Point RECON was **31 psu**, in the optimal range for seagrass, but **above optimal for oysters**.

Water Quality Conditions:

Monitor Site	Salinity (psu) ^a [previous week]	Diss O ₂ (mg/L) ^b	FDOM (qsde) ^c	Chlorophyll (µg/L) ^d	Temperature (°F)
Beautiful Island	1.8 – 3.3 [1.6 – 3.3]	2.8 – 6.1	130	5.0	87.8– 94.4
Fort Myers Yacht Basin	6.3 – 13 [5.6 – 12]	-----	----	----	84.9 – 92.9
Shell Point	24 – 36 [24 – 36]	3.6 –6.9	30	1.5	86.3 – 90.7
McIntyre Creek	31.4 – 36.2 [35.2–36.1]	1.4 – 6.1	13.6 – 36.4	0.9 – 1.7	85.5 – 93.3
Tarpon Bay	32.4 – 36.8 [35.5 – 36.7]	2.9 – 9.1	8.5 – 26.7	0.8 – 3.1	86.1 – 92.2
Wulfert Flats	33.3 – 35.9 [35.0 – 35.8]	0.8 – 7.8	-----	8.4 – 25.3	86.5 – 93.6

Red values are outside of the preferred range.

^a Salinity target values: BI < 5, FM < 10, SP = 10 – 30

^b Dissolved O₂ target values: all sites > 4

^c FDOM target values: BI < 70, FM < 70, SP < 11

^d Chlorophyll target values: BI < 11, FM < 11, SP < 11


^f Temperature target values: < 90

^s Single sonde lower and surface layer or surface grab lab measurement


ND: no data

Wildlife Impacts: In the past week, the CROW wildlife hospital on Sanibel admitted **2 patients** with suspected red tide/toxicosis: 1 adult great blue heron (deceased) and 1 juvenile anhinga (deceased). SCCF documented 1 sea turtle stranding, a deceased Kemp's ridley on Sanibel.

Shellfish Advisory: Shellfish harvest area #6212 (Pine Island Sound Section 1; Aquaculture Lease and Public Reef) is **OPEN** by the Florida Department of Agriculture and Consumer Services (FDACS) as of 6/6/25. SHA #6222 (North Matlacha Pass) and SHA #6232 (South Matlacha Pass) are **OPEN** as of 6/18/25.



USACE Daily Reports			
Date	S79 Flow (cfs)	S78 Flow (cfs)	S77 Flow (cfs)
6/17/25	251	169	0
6/18/25	267	NR	0
6/19/25	278	164	456
6/20/25	419	62	136
6/21/25	356	NR	0
6/22/25	247	99	0
6/23/25	258	146	0
7-day avg	297	128	85



Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (*Vallisneria americana*) health (Ft. Myers only) conditions.

*Ft. Myers sensor is in the lower strata



Water clarity at Lighthouse Beach Park on 6-23-25 at 11:58 AM on a falling tide (3.0 ft).